

2007 International Conference in Honor of Claude Lobry

## Singular perturbations on the infinite time interval

Karim Yadi

Laboratoire des Systèmes Dynamiques et Applications  
Université Aboubekr Belkaïd, BP 119  
13000 Tlemcen, ALGERIE  
yadi@yadikdz@yahoo.fr, k\_yadi@mail.univ-tlemcen.dz

**ABSTRACT.** We consider the slow and fast systems that belong to a small neighborhood of an unperturbed problem. We study the general case where the slow equation has a compact positively invariant subset which is asymptotically stable, and meanwhile the fast equation has asymptotically stable equilibria (Tykhonov's theory) or asymptotically stable periodic orbits (Pontryagin–Rodygin's theory). The description of the solutions is by this way given on infinite time interval. We investigate the stability problems derived from this results by introducing the notion of practical asymptotic stability. We show that some particular subsets of the phase space of the singularly perturbed systems behave like asymptotically stable sets. Our results are formulated in classical mathematics. They are proved within Internal Set Theory which is an axiomatic approach to Nonstandard Analysis.

**RÉSUMÉ.** On considère les systèmes lents-rapides appartenant à un petit voisinage d'un problème non perturbé. On étudie le cas général où l'équation lente admet un sous-ensemble compact positivement invariant qui soit asymptotiquement stable tandis que l'équation rapide a des équilibres asymptotiquement stables (théorie de Tykhonov) ou des cycles limites stables (théorie de Pontryagin). La description des solutions est de ce fait donnée sur des intervalles de temps infinis. On examine les problèmes de stabilité découlant de ces résultats en introduisant la notion de stabilité pratique. On montre que certains sous-ensembles de l'espace de phases des systèmes singulièrement perturbés se comportent comme des ensembles asymptotiquement stables. Les résultats sont formulés classiquement mais sont démontrés dans le cadre de la théorie IST, une approche axiomatique de l'Analyse Non Standard.

**KEYWORDS :** invariant sets, practical asymptotic stability, singular perturbations, nonstandard analysis

**MOTS-CLÉS :** ensembles invariants, stabilité asymptotique pratique, perturbations singulières, analyse non standard

## 1. Introduction

Let us provide the set

$$\mathcal{T} = \{(\Omega, f, g, \alpha, \beta) : \Omega \text{ open subset of } \mathbb{R}^{n+m}, (\alpha, \beta) \in \Omega, \\ f : \Omega \rightarrow \mathbb{R}^n, g : \Omega \rightarrow \mathbb{R}^m \text{ continuous}\}$$

with the topology defined as follows : a neighborhood system of an element

$$(\Omega_0, f_0, g_0, \alpha_0, \beta_0) \in \mathcal{T}$$

is generated by the sets

$$V(D, a) = \{(\Omega, f, g, \alpha, \beta) \in \mathcal{T} : D \subset \Omega, \|f - f_0\|_D < a, \|g - g_0\|_D < a, \\ \|\alpha - \alpha_0\| < a, \|\beta - \beta_0\| < a\},$$

where  $D$  is a compact subset of  $\Omega$  and  $a$  a positive real number. Here,

$$\|h\|_D = \sup_{u \in D} \|h(u)\|,$$

where  $h$  is defined on  $D$  with values in a normed space. Let us call it “the topology of uniform convergence on compacta” and consider the initial value problem

$$\begin{aligned} \varepsilon \dot{x} &= f(x, y), \quad x(0) = \alpha, \\ \dot{y} &= g(x, y), \quad y(0) = \beta, \end{aligned} \tag{1}$$

corresponding to an element  $(\Omega, f, g, \alpha, \beta)$  of  $\mathcal{T}$  where  $\varepsilon$  is a small positive real number and  $(\cdot)' = d/dt$ . The variable  $x$  is called a *fast variable*,  $y$  is called a *slow variable*. We propose to study the system (1) with  $\varepsilon$  small enough and  $(\Omega, f, g, \alpha, \beta)$  sufficiently close to an element  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  of  $\mathcal{T}$  in the sense of the defined topology. It is a singular perturbation problem because the multiplication of the derivative by  $\varepsilon$  fails with the use of theory of continuous dependence of the solutions with respect to the parameters and the initial conditions. Problems as (1) gain their special structure from the presence of two time scales. They are called *slow and fast systems*. We define the *fast equation* by

$$x' = f_0(x, y), \quad y \text{ parameter}, \tag{2}$$

where  $(\cdot)' = d/d\tau$  and  $\tau = t/\varepsilon$ .

Suppose the fast equation admits, for all  $y$  in a compact subset  $Y$  of  $\mathbb{R}^m$  an asymptotically stable equilibrium point  $x = \xi(y)$ , uniformly in the parameter  $y \in Y$ ,  $\xi$  being a continuous function on  $Y$ . We define the *slow equation of* (1) by

$$\dot{y} = g_0(\xi(y), y),$$

and the *reduced problem* by

$$\dot{y} = g_0(\xi(y), y), \quad y(0) = \beta_0. \tag{3}$$

Problem (3) is supposed to have a unique solution  $\bar{y}(t)$  on an interval  $[0, T]$ . It is mainly proved in [13] (see Theorem 4.16 in this paper) that, for  $\varepsilon$  small enough and  $(\Omega, f, g, \alpha, \beta)$

close to  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ , every solution  $(x(t), y(t))$  of (1) is defined at least on  $[0, T]$  and is approximated by  $(\xi(\bar{y}(t)), \bar{y}(t))$  for all  $t \in ]0, T]$ . This approximation can be extended to  $t = 0$  only for  $y(t)$ . Indeed, a boundary layer phenomena is observed for  $x(t)$  at  $t = 0$ . Actually, the fast variable is approximated by the solution of the so-called *boundary layer equation*

$$x' = f_0(x, \beta_0), \quad x(0) = \alpha_0. \quad (4)$$

Now, suppose that the fast equation rather admits for all  $y$  in a compact subset  $G$  of  $\mathbb{R}^m$  a  $T(y)$ -periodic non-trivial solution  $x^*(\tau, y)$ . If the corresponding periodic orbit  $\Gamma_y$  is asymptotically stable, uniformly in  $y \in G$ , we consider the following averaged system as the reduced problem :

$$\dot{y} = \frac{1}{T(y)} \int_0^{T(y)} g_0(x^*(\tau, y), y) d\tau, \quad y(0) = \beta_0. \quad (5)$$

If  $\bar{y}(t)$  is the unique solution of (5), defined on  $[0, T]$ , it is mainly proved in [20] (see Theorem 4.17 in the present paper) that for  $\varepsilon$  small enough and  $(\Omega, f, g, \alpha, \beta)$  close to  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ , every solution  $(x(t), y(t))$  of (1) is defined at least on  $[0, T]$ . The component  $y(t)$  is close to  $\bar{y}(t)$  for  $0 \leq t \leq T$  and the component  $x(t)$  is close to the orbit  $\Gamma_{\bar{y}(t)}$  for  $0 < t \leq T$ . There is a boundary layer at  $t = 0$ . The boundary layer equation (4) describes the fast variable. Hence, after a fast transition, the trajectory rolls up quickly around the manifold generated by the cycles with a slow drift of the  $y$ -component.

For smooth vector fields, when the reduced problem has a limit cycle  $\Gamma$ , the question of whether the singularly perturbed problem has a periodic orbit  $\Gamma_\varepsilon$  near  $\Gamma$  or not for small values of  $\varepsilon$  has been studied by authors like K. O. Friedrichs and W. Wasow [7], L. Flatto and N. Levinson [6], N. D. V. Anosov [1], N. Fenichel [5] and recently F. Verhulst et al. [23, 24]. In [1, 6], it was established that the orbit  $\Gamma$  of period  $\mathcal{P}$  can be continued to a family  $\Gamma_\varepsilon$  of closed orbits if (i)  $\Gamma$ , as an orbit of the reduced problem, has 1 as a simple Floquet multiplier, (ii) for all  $y \in \Gamma$ , the equilibrium point  $x = \xi(y)$  of the boundary layer equation is hyperbolic. This kind of results need strong conditions which can ever ensure both the uniqueness of  $\Gamma_\varepsilon$  for  $\varepsilon$  small enough with a period tending to  $\mathcal{P}$  as  $\varepsilon \rightarrow 0$  and the property of asymptotic phase.

Our aim is first to approximate the solutions of (1) on infinite time interval. More exactly, we will suppose that the slow equation has an asymptotically stable compact subset  $\mathcal{M}$  which is positively invariant. The approximations on unbounded time interval are then given in Section 2 (Theorem 2.3 and Theorem 2.4). If the subset  $\mathcal{M}$  is reduced to an equilibrium point, Theorems 2.3 and 2.4 are nothing else than the extension for all  $t \geq 0$  of Theorems 4.16 and 4.17 already established in [13] and [20] : the solutions of the singularly perturbed system not only live all the time, but they also stay close to the curve formed by the concatenation of the trajectories of the fast and the slow equations. In Section 3, some examples show that there is no hope to obtain stability results with our weak conditions. Nevertheless, the theorems of the preceding section lead to interesting statements in terms of practical stability : when the reduced problem has an asymptotically stable positively invariant compact subset, it arises a subset of the whole phase space which seems to be asymptotically stable when  $\varepsilon \rightarrow 0$  for the singularly perturbed problem (see Theorems 3.2 and 3.3). In the present work, the results are formulated in classical mathematics and proved within *Internal Set Theory* (IST) [15] which is an axiomatic approach of *Nonstandard Analysis* (NSA) [17]. We characterize notions of stability, practical stability and perturbations and translate our main results in nonstandard

words in Section 4. Actually, we rather introduce the notion of s-stability which will lead to results generalizing those of Theorems 3.2 and 3.3. We prove our results in Section 5.

---

## 2. Approximation results

### 2.1. Some definitions

We recall the notion of asymptotic stability (in Lyapunov sense) (see [4, 8] for example) by considering a differential system

$$\dot{x} = f(x), \quad (6)$$

where  $f$  is continuous on an open subset  $U$  of  $\mathbb{R}^n$ .

**Definition 2.1** 1. A subset  $\mathcal{M}$  of  $U$  is said to be **positively invariant** for the system (6) if every solution such that  $x(0) \in \mathcal{M}$  is defined for all  $t > 0$  and satisfies  $x(t) \in \mathcal{M}$ .

2. A bounded subset  $\mathcal{M}$  of  $U$  is said to be **stable** for the system (6) if, for each  $\mu > 0$ , there exists  $\eta > 0$  such that any solution  $x(t)$  of (6) for which  $\text{dis}(x(0), \mathcal{M}) < \eta$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) < \mu$ , where  $\text{dis}(x(t), \mathcal{M})$  denotes the distance from the point  $x(t)$  to the set  $\mathcal{M}$  given by  $\inf_{m \in \mathcal{M}} \|x(t) - m\|$ .

3.  $\mathcal{M}$  is said to be **attractive** if it admits a neighborhood  $\mathcal{V}$  (basin of attraction) such that any solution  $x(t)$  of (6) for which  $x(0) \in \mathcal{V}$  can be continued for all  $t \geq 0$  and satisfies  $\lim_{t \rightarrow \infty} \text{dis}(x(t), \mathcal{M}) = 0$ .

3.  $\mathcal{M}$  is **asymptotically stable** if it is stable and attractive.

Consider the case where the system

$$\dot{x} = f(x, y), \quad (7)$$

depends on a parameter  $y$  which belongs to a set  $Y$ . Let  $\mathcal{M}_y$  be an asymptotically stable bounded subset of  $U$  for each value of  $y$  in  $Y$ .

**Definition 2.2** The basin of attraction of the asymptotically stable bounded subset  $\mathcal{M}_y$  of  $U$  is said to be **uniform over  $Y$**  for (7) if there exists a real number  $a > 0$  such that, for all  $y$  in  $Y$ , the set  $\{x \in \mathbb{R}^n : \text{dis}(x, \mathcal{M}_y) \leq a\}$  is in the basin of attraction of  $\mathcal{M}_y$ .

### 2.2. The main theorems

Let us first make the following assumptions :

*T1* : The fast equation (2) has the property of uniqueness of the solutions with the prescribed initial conditions.

The slow manifold of the system (1) is defined by the set of the points of  $\mathbb{R}^n \times \mathbb{R}^m$  such that

$$f_0(x, y) = 0. \quad (8)$$

It is namely the set of equilibria of the fast equation (2).

*T2* : There exists a continuous mapping  $\xi : Y \rightarrow \mathbb{R}^n$ ,  $Y$  being a compact subset of  $\mathbb{R}^m$  with a non empty interior, such that  $(\xi(y), y) \in \Omega_0$  for all  $y \in Y$  and  $f_0(\xi(y), y) = 0$ . Moreover, for all  $y \in Y$ ,  $x = \xi(y)$  is an isolated root of the equation (8), that is there exists a real number  $\delta > 0$  such that if  $y \in Y$ ,  $\|x - \xi(y)\| < \delta$  and  $x \neq \xi(y)$  then  $f_0(x, y) \neq 0$ .

The subset  $\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^m : x = \xi(y), y \in Y\}$  of  $\mathbb{R}^n \times \mathbb{R}^m$  is an  $m$ -dimensional manifold included in the slow manifold.

*T3* : For all  $y$  in  $Y$ , the equilibrium point  $x = \xi(y)$  of the fast equation (2) is asymptotically stable and its basin of attraction is uniform over  $Y$ .

By substituting  $\xi(y)$  to  $x$  in the second equation of (1) we obtain the slow equation

$$\dot{y} = g_0(\xi(y), y), \quad (9)$$

which will be defined in the interior  $\overset{\circ}{Y}$  of the compact set  $Y$ .

*T4* : The slow equation (9) has the property of uniqueness of the solutions with the prescribed initial conditions.

*T5* : The point  $\beta_0$  is in  $\overset{\circ}{Y}$  and  $\alpha_0$  is in the basin of attraction of the equilibrium point  $x = \xi(\beta_0)$ .

**Theorem 2.3** Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  be an element of  $\mathcal{T}$  and  $\xi : Y \rightarrow \mathbb{R}^n$  be a continuous function. Let hypotheses *T1* to *T5* be satisfied. Let  $\mathcal{M}$  be a closed subset in  $\overset{\circ}{Y}$  which is positively invariant for the slow equation (9). Suppose that  $\mathcal{M}$  is asymptotically stable for (9) with  $\beta_0$  in its basin of attraction. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (3). Then, for all  $\eta > 0$ , there exists a real number  $\varepsilon^* > 0$  and a neighborhood  $\mathcal{V}$  of the element  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  of  $\mathcal{T}$  such that, for all  $\varepsilon < \varepsilon^*$  and all  $(\Omega, f, g, \alpha, \beta) \in \mathcal{V}$ , any solution  $(x(t), y(t))$  of the problem (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that :

$$\begin{aligned} \varepsilon\omega &< \eta, \quad 1/\omega' < \eta, \\ \|x(\varepsilon\tau) - \tilde{x}(\tau)\| &< \eta \text{ for } 0 \leq \tau \leq \omega, \\ \|x(t) - \xi(\bar{y}(t))\| &< \eta \text{ for } t \in [\varepsilon\omega, \omega'] \\ \|y(t) - \bar{y}(t)\| &< \eta \text{ for } t \in [0, \omega'], \\ \text{dis}(y(t), \mathcal{M}) &< \eta \text{ for } t \geq \omega', \\ \text{dis}(x(t), \xi(\mathcal{M})) &< \eta \text{ for } t \geq \omega'. \end{aligned} \quad (10)$$

The set  $\xi(\mathcal{M})$  is the range of  $\mathcal{M}$  by the function  $\xi$ . Note that  $\mathcal{M}$  is necessarily a compact subset of  $\overset{\circ}{Y}$  as will be all the positively invariant subsets of the following. When the fast equation has limit cycles, the preceding theorem does not fit. Let us make instead the following assumptions, the first one being nothing else than (*T1*).

*P1* : The fast equation (2) has the property of uniqueness of the solutions with the prescribed initial conditions.

*P2* : There exists a family of solutions  $x^*(\tau, y)$  depending continuously on  $y \in G$ , where  $G$  is a compact subset of  $\mathbb{R}^m$  with a non empty interior, such that  $x^*(\tau, y)$  is a periodic solution of the fast equation (2) of period  $T(y) > 0$ , the mapping  $y \rightarrow T(y)$  is continuous.

*P3* : The closed orbit  $\Gamma_y$  corresponding to the periodic solution  $x^*(\tau, y)$  is asymptotically stable and its basin of attraction is uniform over  $G$ .

What precede means that the cycle<sup>1</sup>  $\Gamma_y$  depends continuously on  $y$  and is locally unique, that is, there exists an neighborhood  $W$  of  $\Gamma_y$  such that the equation (2) has no other cycle in  $W$ .

We define the slow equation in the interior  $\mathring{G}$  of  $G$  by the averaged system

$$\dot{y} = \bar{g}_0(y) := \frac{1}{T(y)} \int_0^{T(y)} g_0(x^*(\tau, y), y) d\tau. \tag{11}$$

Assume what follows :

*P4* : The slow equation (11) has the uniqueness of the solutions with the prescribed initial conditions.

*P5* : The point  $\beta_0$  is in  $\mathring{G}$  and  $\alpha_0$  is in the basin of attraction of  $\Gamma_{\beta_0}$ .

**Theorem 2.4** *Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  be an element of  $\mathcal{T}$ . Let hypotheses P1 to P5 be satisfied. Let  $\mathcal{M}$  be a closed subset in  $\mathring{G}$  which is positively invariant for the slow equation (11). Suppose that  $\mathcal{M}$  is asymptotically stable for (11) with  $\beta_0$  in its basin of attraction. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (5). Then, for all  $\eta > 0$ , there exists a real number  $\varepsilon^* > 0$  and a neighborhood  $\mathcal{V}$  of the element  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  of  $\mathcal{T}$  such that, for all  $\varepsilon < \varepsilon^*$  and all  $(\Omega, f, g, \alpha, \beta) \in \mathcal{V}$ , any solution  $(x(t), y(t))$  of the problem (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that :*

$$\begin{aligned} \varepsilon\omega &< \eta, \quad 1/\omega' < \eta, \\ \|x(\varepsilon\tau) - \tilde{x}(\tau)\| &< \eta \text{ for } 0 \leq \tau \leq \omega, \\ \|y(t) - \bar{y}(t)\| &< \eta \text{ for } t \in [0, \omega'], \\ \text{dis}(x(t), \Gamma_{\bar{y}(t)}) &< \eta \text{ for } t \in [\varepsilon\omega, \omega'], \\ \text{dis}(y(t), \mathcal{M}) &< \eta \text{ for } t \geq \omega', \\ \text{dis}(x(t), \Gamma_{y(t)}) &< \eta \text{ for } t \geq \omega'. \end{aligned}$$

The proofs of Theorems 2.3 and 2.4 are postponed to Section 4.5. The particular case of Theorem 2.3, when  $\mathcal{M}$  is an asymptotically stable equilibrium point, is Theorem 2 of [13]. The result therein concerns the case where the slow equation (9) has an asymptotically stable equilibrium point  $y_\infty$  in  $\mathring{Y}$  with  $\beta_0$  in its basin of attraction. For  $\varepsilon$  small enough and for  $(\Omega, f, g, \alpha, \beta)$  close to  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ , under the assumptions T1 to T5, any solution  $(x(t), y(t))$  of (1) is close to  $(\xi(\bar{y}(t)), \bar{y}(t))$  for all  $t > 0$ . Here, one takes  $\mathcal{M} = \{y_\infty\}$ . The particular case of Theorem 2.4, when  $\mathcal{M}$  is an asymptotically stable equilibrium point, is Theorem 2.3 of [20]. Under the assumptions P1 to P5, any solution  $(x(t), y(t))$  of (1) is close to  $(\Gamma_{\bar{y}(t)}, \bar{y}(t))$  for all  $t > 0$ . Finally, one can mention the important particular cases where the slow equations (9) and (11) admit an asymptotically stable cycle.

---

1. Actually, the term cycle should be used for planar systems. A stable limit cycle is the asymptotically stable closed orbit associated to an orbitally asymptotically stable periodic solution.

### 3. Stability results

#### 3.1. Some examples

Suppose that the slow equation (9) admits, say an asymptotically stable equilibrium point  $y_\infty$  and that the corresponding point  $(\xi(y_\infty), y_\infty)$  lying on the slow manifold is an equilibrium point of (1). To deduce asymptotic stability of  $(\xi(y_\infty), y_\infty)$  for the whole system (1) for  $\varepsilon$  small enough, Tykhonov theory needs strong assumptions like smoothness of the field and exponential stability of the equilibrium points of both slow and fast equations (see [9], Theorem 9.3, page 380<sup>2</sup>). It is also the case in Pontryagin-Rodygin theory. In most of the applications, it is quite reasonable to impose conditions of exponential or uniform exponential stability instead of the simple asymptotic stability (asymptotic stability which is not exponential is said to be critical). The following examples show the non robustness of the critical asymptotic stability. Before, one has to keep in mind that the condition of exponential stability for such results is sufficient, but not necessary as shown in the following example (see also [9], Theorem 9.2, page 377).

**Example 1** ([9], Example 9.9, page 377) The origin of the system

$$\begin{aligned}\varepsilon \dot{x} &= -x + \varepsilon(x - y^3), \\ \dot{y} &= -y^3 + x.\end{aligned}\tag{12}$$

is an equilibrium point. The slow equation is give by

$$\dot{y} = -y^3.$$

Its origin is asymptotically but not exponentially stable. The fast equation is

$$x' = -x,$$

and its origin is exponentially stable. It is shown in [9] that for  $\varepsilon$  small enough, the origin of the the whole system (12) is asymptotically stable by the use of the so-called quadratic Lyapounov functions. It is well explained in the same reference that the existence of quadratic Lyapounov functions do not always imply exponential stability which is a particular case.

**Example 2** The fast equation of the planar slow and fast system

$$\begin{aligned}\varepsilon \dot{x} &= -x + y, \\ \dot{y} &= -y^2 x + \varepsilon y,\end{aligned}\tag{13}$$

where  $\varepsilon > 0$  is a small real parameter, is given by

$$x' = -x + y.$$

Assumption *T1* is clearly satisfied for the fast equation. The slow curve is the line  $x = \xi(y) := y$  (Assumption *T2*). All it points are (globally) asymptotically stable equilibria

---

2. Actually, the theorem therein concerns the nonautonomous case but can be stated or the autonomous one.

of the fast equation for each value of  $y$  (Assumption T3). By substituting  $\xi(y)$  to  $x$  and 0 to  $\varepsilon$  in the second equation of the system, we obtain the slow equation

$$\dot{y} = -y^3,$$

which satisfies Assumption T4. The origin  $y = 0$  of the slow equation is (globally) asymptotically stable (Assumption T6). One can apply the preceding theory (more exactly the extension of Tykhonov's Theorem for infinite time interval) for any initial condition  $(x_0, y_0) \neq (0, 0)$  (Assumption T5) to have

$$\begin{aligned} \forall t \geq 0, \lim_{\varepsilon \rightarrow 0} y(t, \varepsilon) &= \bar{y}(t), \\ \forall t > 0, \lim_{\varepsilon \rightarrow 0} x(t, \varepsilon) &= \xi(\bar{y}(t)) = \bar{y}(t), \end{aligned}$$

where  $(x(t, \varepsilon), y(t, \varepsilon))$  is the solution of the whole system with initial condition  $(x_0, y_0)$  and  $\bar{y}(t)$  is the solution of the reduced problem. Now, knowing that  $\bar{y}(t)$  goes to 0 when  $t$  goes towards  $+\infty$ , we obtain that

$$\lim_{\varepsilon \rightarrow 0, t \rightarrow +\infty} (x(t, \varepsilon), y(t, \varepsilon)) = (0, 0).$$

However, this limit does not mean that the origin of the whole system, which is an equilibrium point, is asymptotically stable, not even attractive. It is easy to check, by linearization, that it is a saddle point for any positive value of  $\varepsilon$ .

**Example 3** The following example

$$\begin{aligned} \varepsilon \dot{x} &= -x^3 + \varepsilon x, \\ \dot{y} &= -y + x^2, \end{aligned} \tag{14}$$

is given just to exhibit a case where the origin of the fast equation  $x' = -x^3$  is critically asymptotically stable whereas the origin of the slow equation  $\dot{y} = -y$  is exponentially stable. One can claim that any solution of the singularly perturbed system goes towards the equilibrium point  $(0, 0)$  when  $\varepsilon \rightarrow 0$  and  $t \rightarrow +\infty$ . Nevertheless, the origin  $(0, 0)$  is a saddle point for each value of  $\varepsilon > 0$ .

**Example 4** The fast and the slow equation of the system

$$\begin{aligned} \varepsilon \dot{x} &= -x + \varepsilon, \\ \dot{y} &= -y + \varepsilon x, \end{aligned}$$

are respectively given by  $x' = -x$  and  $\dot{y} = -y$ . In both cases, the origin is an exponentially stable equilibrium point. Any solution of the singularly perturbed system goes towards the equilibrium point  $(0, 0)$  when  $\varepsilon \rightarrow 0$  and  $t \rightarrow +\infty$ . Nevertheless, the origin  $(0, 0)$  is not even an equilibrium point of the whole system.

**Example 5** The following tridimensional singularly perturbed system  $(\Sigma)$

$$\begin{aligned} \varepsilon \dot{x}_1 &= -x_2 + x_1 \left( \sqrt{x_1^2 + x_2^2} - 1 + \varepsilon \right) \left( \sqrt{x_1^2 + x_2^2} - 1 - \varepsilon \right) (1 - x_1^2 - x_2^2), \\ \varepsilon \dot{x}_2 &= x_1 + x_2 \left( \sqrt{x_1^2 + x_2^2} - 1 + \varepsilon \right) \left( \sqrt{x_1^2 + x_2^2} - 1 - \varepsilon \right) (1 - x_1^2 - x_2^2), \\ \dot{y} &= -y^3 x_1^2, \end{aligned}$$

is written under cylindrical coordinates  $(x_1 = r \cos \theta, x_2 = r \sin \theta, y)$  as

$$\begin{aligned}\varepsilon \dot{r} &= r(r-1+\varepsilon)(r-1-\varepsilon)(1-r^2), \\ \varepsilon \dot{\theta} &= 1, \\ \dot{y} &= -r^2 y^3 \cos^2 \theta.\end{aligned}$$

The equilibrium point  $r = 1$  of the first equation of

$$\begin{aligned}r' &= r(r-1)^2(1-r^2), \\ \theta' &= 1,\end{aligned}\tag{15}$$

corresponds to a stable limit cycle  $\Gamma_y$  the basin of attraction of which is the whole plane  $(x_1, x_2)$  except the origin. So, it is uniform in  $y$  (actually the cycle does not depend on  $y$ ). A  $2\pi$ -periodic solution associated to the cycle is  $x^*(\tau, y) = (\cos \tau, \sin \tau)$ . According to Pontryagin-Rodygin's theory, the equation describing the slow motion is given by the average system

$$\dot{y} = -\frac{1}{2\pi} \int_0^{2\pi} y^3 \cos^2 \tau d\tau = -\frac{y^3}{2},\tag{16}$$

the origin  $y = 0$  of which is an asymptotically stable equilibrium. The preceding theory, and more exactly the extension of Pontryagin-Rodygin's Theorem in [20], implies that for any initial condition  $(x_1^0, x_2^0, y^0)$  of  $(\Sigma)$ , apart from the origin, we have

$$\forall t \geq 0, \lim_{\varepsilon \rightarrow 0} y(t, \varepsilon) = \bar{y}(t), \quad \forall t > 0, \lim_{\varepsilon \rightarrow 0} \text{dis}((x_1(t, \varepsilon), x_2(t, \varepsilon)), \Gamma_{\bar{y}(t)}) = 0.$$

Here,  $(x_1(t, \varepsilon), x_2(t, \varepsilon), y(t, \varepsilon))$  is the solution of the singularly perturbed problem  $(\Sigma)$  with initial condition  $(x_1^0, x_2^0, y^0)$  and  $\bar{y}(t)$  is the solution of (16) starting from  $y^0$ . Since  $\bar{y}(t) \rightarrow 0$  when  $t \rightarrow +\infty$ , we have the following limit

$$\lim_{\varepsilon \rightarrow 0, t \rightarrow +\infty} \text{dis}((x_1(t, \varepsilon), x_2(t, \varepsilon), y(t, \varepsilon)), \Gamma_0 \times \{0\}) = 0.$$

This does not either mean that the closed curve  $\Gamma_0 \times \{0\}$  in the phase space  $\mathbb{R}^3$  is an asymptotically stable orbit. One can see that for each  $\varepsilon > 0$ , the cylinder generated by the cycles  $\Gamma_y$  is "repelling" for  $(\Sigma)$ , being located between two "attracting" cylinders corresponding to  $r = 1 - \varepsilon$  and  $r = 1 + \varepsilon$ . In other words, the "exact fast dynamics" of the whole problem, that is the subsystem

$$\begin{aligned}r' &= r(r-1+\varepsilon)(r-1-\varepsilon)(1-r^2), \\ \theta' &= 1,\end{aligned}$$

which is a regular perturbation of (15), admits for all  $y$  and all  $\varepsilon > 0$  two "stable limit cycles"  $r = 1 - \varepsilon$  and  $r = 1 + \varepsilon$  surrounding the "unstable limit cycle"  $r = 1$ .

**Example 6** The slow dynamic associated to the following slow-fast system written in cylindrical coordinates  $(x = x, y_1 = r \cos \theta, y_2 = r \sin \theta)$

$$\begin{aligned}\varepsilon \dot{x} &= -x + r \cos \theta, \\ \dot{r} &= r(r-1+\varepsilon)(r-1-\varepsilon)(1-r^2), \\ \dot{\theta} &= 1,\end{aligned}\tag{17}$$

is approximated by the slow equation

$$\begin{aligned}\dot{r} &= -r(r-1)^3(r+1), \\ \dot{\theta} &= 1,\end{aligned}$$

having the unit circle  $\Gamma$  as a unique asymptotically stable cycle. The equilibrium points of the fast equation  $x' = -x + y_1$  are all points of the slow surface  $x = \xi(y_1, y_2) := y_1$  and they are (globally) asymptotically stable. Let  $(x(t, \varepsilon), y_1(t, \varepsilon), y_2(t, \varepsilon))$  be the solution of (17) written in the rectangular coordinates with any initial condition apart the origin. We can deduce from Theorem 2.3 in the special case where  $\mathcal{M} = \Gamma$  that the following limits hold

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0, t \rightarrow +\infty} \text{dis}(x(t, \varepsilon), \xi(\Gamma)) &= 0, \\ \lim_{\varepsilon \rightarrow 0, t \rightarrow +\infty} \text{dis}((y_1(t, \varepsilon), y_2(t, \varepsilon)), \Gamma) &= 0. \end{aligned}$$

But even if the subset  $\xi(\Gamma) \times \Gamma$  of  $\mathbb{R}^3$  was properly a closed orbit of (17), it could not be stable. Indeed, for the subsystem

$$\begin{aligned} \dot{r} &= r(r - 1 + \varepsilon)(r - 1 - \varepsilon)(1 - r^2), \\ \dot{\theta} &= 1, \end{aligned}$$

of (17), for any value of  $\varepsilon > 0$ , the orbit  $\Gamma$  is an unstable limit cycle, surrounded by the two stable limit cycles  $r = 1 - \varepsilon$  and  $r = 1 + \varepsilon$ .

### 3.2. Practical stability

Although they do not imply the existence of asymptotically stable sets for the singularly perturbed systems, the limits of the preceding examples show a kind of “seeming” stability of certain points or closed curves in the whole phase space. It is known (see [8], Theorem 38.1) that the uniform attractivity, say of the origin of a system, with respect to the initial conditions, implies its asymptotic stability. Moreover, the global asymptotic stability is equivalent to the uniformity of the attractivity for any initial condition in an arbitrarily large ball centered at the origin<sup>3</sup>. The following definition is inspired by lectures given by Lobry and Sari during the CIMPA School-2003 in Tlemcen, *Contrôle Non Linéaire et Applications* [12] (see also [11]). The terminology is yet taken from control theory where it arises in problems of stabilization (see for example [21]). Let us consider the slow and fast system

$$\begin{aligned} \varepsilon \dot{x} &= f_0(x, y), \\ \dot{y} &= g_0(x, y), \end{aligned} \tag{18}$$

where  $f_0 : \Omega_0 \rightarrow \mathbb{R}^n$  and  $g_0 : \Omega_0 \rightarrow \mathbb{R}^m$  are continuous on an open subset  $\Omega_0$  of  $\mathbb{R}^{n+m}$ .

**Definition 3.1** *A bounded subset  $\mathcal{A}$  of  $\Omega_0$  is said to be semiglobally practically asymptotically stable (SGPAS) for the system (18) when  $\varepsilon \rightarrow 0$  if, for every compact neighborhood  $\mathcal{K} \subset \Omega_0$  of  $\mathcal{A}$  and every neighborhood  $\mathcal{O} \subset \Omega_0$  of  $\mathcal{A}$ , there exist  $\varepsilon_0 > 0$  and a real number  $T > 0$  such that, for all  $\varepsilon \in ]0, \varepsilon_0]$ , all  $t \geq T$  and all  $(\alpha, \beta) \in \mathcal{K}$ , any solution  $(x(t), y(t))$  of (18) with initial condition  $(\alpha, \beta)$  is in  $\mathcal{O}$ .*

*If  $\mathcal{K}$  is not arbitrary, the subset  $\mathcal{A}$  is said to be practically asymptotically stable (PAS) for the system (18) when  $\varepsilon \rightarrow 0$ .*

The definition says that any solution  $(x(t), y(t))$  of (18) with initial condition  $(\alpha, \beta)$  satisfies  $\lim_{t \rightarrow +\infty} \text{dis}((x(t), y(t)), \mathcal{A}) = 0$  when  $t \rightarrow +\infty$  and  $\varepsilon \rightarrow 0$ , the convergence being

3. The famous Vinograd example ([?], page 191) is a good illustration of attractivity which is not uniform with respect to the initial conditions.

uniform with respect to  $(\alpha, \beta)$  in  $\mathcal{K}$  (i.e.  $\lim_{t \rightarrow +\infty, \varepsilon \rightarrow 0} \text{Sup}_{(\alpha, \beta) \in \mathcal{K}} \text{dis}((x(t), y(t)), \mathcal{A}) = 0$ ). In other words, for the first part of the definition, any trajectory starting from an arbitrarily point  $(\alpha, \beta)$  which is not close to the boundary of  $\Omega_0$  reaches an arbitrarily small neighborhood  $\mathcal{O}$  of  $\mathcal{A}$  in finite time and for  $\varepsilon$  small enough. Note that it is required that the finite time  $T$  and the threshold  $\varepsilon_0$  are the same for any initial condition  $(\alpha, \beta)$ .

### 3.3. The theorems


Finally, practical stability answers to natural questions as the following : “If the slow equation (9) admits, say an asymptotically stable equilibrium point  $y_\infty$  or a limit cycle  $\Gamma$ , which part acts, for the whole problem (1), the corresponding point  $(\xi(y_\infty), y_\infty)$  or subset  $\xi(\Gamma) \times \Gamma$  lying on the slow manifold? They have no reason to be respectively an equilibrium point or a cycle, unless we add strong hypotheses like smoothness and exponential stability. However, for very small values of the parameter  $\varepsilon$  they look like. Because of the asymptotic feature of this property, an engineer, a biologist,...would be satisfied by a model which seems to tend towards a steady or permanently oscillating state, instead of a perfect stability which is after all ideal. The following results of practical stability deduce from the Theorems 2.3 and 2.4. Their proofs are postponed to Section 4.5.

**Theorem 3.2** *Let  $f_0 : \Omega_0 \rightarrow \mathbb{R}^n$  and  $g_0 : \Omega_0 \rightarrow \mathbb{R}^m$  be continuous on an open subset  $\Omega_0$  of  $\mathbb{R}^{n+m}$  satisfying hypotheses T1 to T4. Let  $\mathcal{M}$  be a closed subset in  $\bar{Y}$  which is positively invariant for the slow equation (9). Suppose that  $\mathcal{M}$  is asymptotically stable for (9). Then the subset  $\xi(\mathcal{M}) \times \mathcal{M}$  of the slow manifold is PAS for the system (18) when  $\varepsilon \rightarrow 0$ .*


**Theorem 3.3** *Let  $f_0 : \Omega_0 \rightarrow \mathbb{R}^n$  and  $g_0 : \Omega_0 \rightarrow \mathbb{R}^m$  be continuous on an open subset  $\Omega_0$  of  $\mathbb{R}^{n+m}$  satisfying hypotheses P1 to P4. Let  $\mathcal{M}$  be a closed subset in  $\bar{G}$  which is positively invariant for the slow equation (11). Suppose that  $\mathcal{M}$  is asymptotically stable for (11). Then the subset  $\bigcup_{y \in \mathcal{M}} (\Gamma_y \times \{y\})$  of  $\Omega_0$  is PAS for the system (18) when  $\varepsilon \rightarrow 0$ .*

If the equilibrium points and the positively invariant sets of the fast and slow equations were globally asymptotically stable, we would have results in terms of SGPAS when  $\varepsilon \rightarrow 0$ . A natural consequence of these two theorems is obtained when  $\mathcal{M}$  is an asymptotically stable cycle. In the case of Theorem 3.2, the singularly perturbed system (18) seems to have an asymptotically stable cycle on the slow manifold for  $\varepsilon$  small enough. In the case of Theorem 3.3, we obtain a torus which seems to attract the solutions of (18) when  $\varepsilon$  is sufficiently small. Hence, in the light of these results, we can come back to the examples above. In Example 2, the equilibria of the fast equation are exponentially stable, the origin of the slow equation is asymptotically stable, the origin is an equilibrium point of the whole system and it is SGPAS when  $\varepsilon \rightarrow 0$ . In Example 3, the origin is asymptotically stable for the fast equation, exponentially stable for the slow equation, it is an equilibrium point of the whole system and this equilibrium is SGPAS when  $\varepsilon \rightarrow 0$ . In Example 4, the origin of both the fast and the slow equation is exponentially stable, but  $(0, 0)$  is not an equilibrium point of the singularly perturbed system. However, it is SGPAS when  $\varepsilon \rightarrow 0$ . In Example 5, the fast equation has a unique asymptotically stable cycle  $\Gamma_y$  for each value of  $y$ , the origin of the slow equation is asymptotically stable. The subset  $\Gamma_0 \times \{0\}$  of  $\mathbb{R}^3$  is SGPAS when  $\varepsilon \rightarrow 0$  (even if it was a cycle, it would not be stable). Finally, in the Example 6, the slow equation has a unique asymptotically stable cycle  $\Gamma$ , the equilibrium

of the fast equation is exponentially stable for each  $(y_1, y_2)$ . The subset  $\xi(\Gamma) \times \Gamma$  of  $\mathbb{R}^3$  is SGPAS when  $\varepsilon \rightarrow 0$ . As an illustration, the numerical simulations of system  $(\Sigma)$  in example 5 with Mapple show that the more  $\varepsilon$  is small, the more the trajectories seem to tend to the cycle  $\Gamma_0 \times \{0\}$  although it is unstable.



**Figure 1.** Numerical simulation of Example 4 with  $\varepsilon = 0.1$ .



**Figure 2.** Numerical simulation of Example 4 with  $\varepsilon = 0.003$ .

---

#### 4. External formulations

The Internal Set Theory is an extension of ordinary mathematics. It gives an axiomatic approach to A. Robinson's Nonstandard Analysis [17] and has been elaborated by E. Nelson [15]. Tutorial on IST is available in many articles as for example in [13, 20]. For interested readers, *Nonstandard Analysis in Practice* [3] is a recommended book deal-

ing with at least nine fields where non standard analysis has been used. One can read about the perturbation theory of ordinary differential equations in [2, 10, 14, 18, 19, 25]. To the binary undefined predicate  $\in$  of the classical Set Theory ZFC (Zermelo-Fraenkel plus axiom of choice), a new unary undefined predicate *standard* (st) is joined. The new predicate is governed by three new axioms added to ZFC. A formula of IST is said to be internal in the case where it does not involve the new predicate st. Otherwise, it is called external. Some theorems proved in IST are external and can be reformulated so they become internal. Indeed, there is a *reduction algorithm* due to Nelson which reduces any external formula  $F(x_1, \dots, x_n)$  of IST without other free variables than  $x_1, \dots, x_n$  to an internal formula  $F'(x_1, \dots, x_n)$  with the same free variables, such that  $F \equiv F'$ , that is,  $F \Leftrightarrow F'$  for all standard values of the free variables. We will need the following reduction formulas which occur frequently

$$\begin{aligned} \forall x (\forall^{\text{st}} y A \Rightarrow \forall^{\text{st}} z B) &\equiv \forall z \exists^{\text{fin}} y' \forall x (\forall y \in y' A \Rightarrow B), \\ \forall x (\exists^{\text{st}} w \forall^{\text{st}} y A \Rightarrow \forall^{\text{st}} z B) &\equiv \forall w \forall z \exists^{\text{fin}} y' \forall x (\forall y \in y' A \Rightarrow B), \end{aligned} \quad (19)$$

where  $A$  (respectively  $B$ ) is an internal formula with free variable  $y$  (respectively  $z$ ) and standard parameters. The notation  $\forall^{\text{st}}$  means “for all standard” and  $\exists^{\text{fin}}$  means “there is a finite”.

#### 4.1. The short shadow lemma

We recall here an important result of non standard analysis which generalizes the classical theory of dependence of solutions with respect to parameters and initial conditions. The statement of the so-called *Short Shadow Lemma* we give here is more general than the version we will consider in this paper (Lemma 4.3). Nevertheless, this general version was used in the proof of the essential results of Theorems 4.16 and 4.17. This efficient tool concerns the theory of regular perturbations. Consider the following initial value problems :

$$\frac{dx}{dt} = F_0(x), \quad x(0) = a_0 \in U_0, \quad (20)$$

$$\frac{dx}{dt} = F(x), \quad x(0) = a \in U. \quad (21)$$

The Short Shadow Lemma permits the comparison of the solutions of (20) and (21) when  $F$  is close to  $F_0$  and  $a$  is close to  $a_0$  in a certain meaning. One can find a proof in [19] with the use of an other powerful result, the *Stroboscopy Lemma*. This last was the most important tool in the proof of Theorem 4.17 in [20].

**Theorem 4.1 (Short Shadow Lemma)** *Let  $U_0$  a standard open subset of  $\mathbb{R}^n$  and let  $F_0 : U_0 \rightarrow \mathbb{R}^n$  be standard and continuous. Let  $a_0 \in U_0$  be standard. Suppose that the initial value problem (20) has a unique solution  $x_0(t)$  and let  $J = [0, \omega[$ ,  $0 < \omega \leq +\infty$ , its maximal positive interval of definition. Let  $U$  an open subset of  $\mathbb{R}^n$  containing all the nearstandard elements in  $U_0$ . Let  $F : U \rightarrow \mathbb{R}^n$  be continuous such that  $F(x) \simeq F_0(x)$  for all  $x$  nearstandard in  $U_0$ . Then, every solution  $x(t)$  of the initial value problem (21) with  $a \simeq a_0$  is defined for all  $t$  nearstandard in  $J$  and satisfies  $x(t) \simeq x_0(t)$ .*

## 4.2. External characterization of stability

The external formulation of stability of a bounded subset  $\mathcal{M} \subset U$  of the system (6) is given by the following lemma.

**Lemma 4.2** *Assume that  $f, U$  and  $\mathcal{M}$  are standard. The subset  $\mathcal{M} \subset U$  is :*

1. *Stable for the system (6), if and only if any solution  $x(t)$  of (6) for which  $\text{dis}(x(0), \mathcal{M}) \simeq 0$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$ .*
2. *Attractive if and only if  $\mathcal{M}$  admits a standard neighborhood  $\mathcal{V}$  (basin of attraction) such that any solution  $x(t)$  of (6) for which  $x(0)$  is standard in  $\mathcal{V}$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all  $t \simeq +\infty$ .*

**Proof.** 1. Let us denote by  $B$  the formula “any solution  $x(t)$  of (6) for which  $x(0) = \alpha$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) < \mu$ .” To say in the lemma  $\text{dis}(\alpha, \mathcal{M}) \simeq 0$  is the same as to say  $\forall^{\text{st}} \eta \text{dis}(\alpha, \mathcal{M}) < \eta$  and to say  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  is the same as to say  $\forall^{\text{st}} \mu \text{dis}(x(t), \mathcal{M}) < \mu$ . The stability of  $\mathcal{M}$  is then characterized by the formula

$$\forall \alpha (\forall^{\text{st}} \eta \text{dis}(\alpha, \mathcal{M}) < \eta \Rightarrow \forall^{\text{st}} \mu B),$$

where  $f$  and  $\mathcal{M}$  are standard parameters and  $\eta, \mu$  range over the positive real numbers. By the first reduction formula (19), this is equivalent to

$$\forall \mu \exists^{\text{fin}} \eta' \forall \alpha (\forall \eta \in \eta' \text{dis}(\alpha, \mathcal{M}) < \eta \Rightarrow B).$$

The set  $\eta'$  being finite, there exists  $\eta$  such that  $\eta = \min \eta'$  and the last formula becomes

$$\forall \mu \exists \eta \forall \alpha (\text{dis}(\alpha, \mathcal{M}) < \eta \Rightarrow B),$$

which is nothing else than the usual definition of stability of  $\mathcal{M}$ . 2. By transfer, the attractivity of a standard subset is equivalent to the existence of a standard basin of attraction. In the lemma, to say  $t \simeq +\infty$  is the same as to say  $\forall^{\text{st}} r t > r$ . Thus the characterization of the standard basin of attraction  $\mathcal{V}$  is that any solution  $x(t)$  of the equation (6) for which  $x(0)$  is standard in  $\mathcal{V}$  can be continued for all  $t \geq 0$  and satisfies

$$\forall t (\forall^{\text{st}} r t > r \Rightarrow \forall^{\text{st}} \mu \text{dis}(x(t), \mathcal{M}) < \mu).$$

In this formula,  $x(\cdot)$  and  $\mathcal{M}$  are standard parameters while  $r, \mu$  range over the positive real numbers. By (19), this is equivalent to

$$\forall \mu \exists^{\text{fin}} r' \forall t (\forall r \in r' t > r \Rightarrow \text{dis}(x(t), \mathcal{M}) < \mu).$$

To say, for  $r'$  a finite set,  $\forall r \in r' t > r$  is the same as to say  $t > r$  for  $r = \max r'$  and the formula is equivalent to

$$\forall \mu \exists r \forall t (t > r \Rightarrow \text{dis}(x(t), \mathcal{M}) < \mu).$$

Finally, for all standard  $\alpha$  in  $\mathcal{V}$ , any solution  $x(t)$  of the equation (6) for which  $x(0) = \alpha$ , can be continued for all  $t \geq 0$  and satisfies  $\lim_{t \rightarrow \infty} \text{dis}(x(t), \mathcal{M}) = 0$ . By transfer, this property remains true for all  $\alpha$  in  $\mathcal{V}$ . This is the usual definition of the attractivity. ■

Here is an elegant characterization of the asymptotic stability of a compact positively invariant subset.

**Lemma 4.3** *Assume that the system (6) has the property of uniqueness of the solutions with the prescribed initial conditions and that  $f$ ,  $U$  and  $\mathcal{M}$  are standard. Assume that  $\mathcal{M}$  is a compact subset of  $U$  positively invariant for (6). Then  $\mathcal{M}$  is asymptotically stable for the system (6) if and only if there exists a standard  $a > 0$  such that any solution  $x(t)$  of (6) for which  $\text{dis}(x(0), \mathcal{M}) < a$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all  $t \simeq +\infty$ .*

**Proof.** Assume that  $\mathcal{M}$  is asymptotically stable. By attractivity, it has a standard basin of attraction  $\mathcal{V}$ . Let  $a > 0$  be standard such that the closure of the set  $A = \{x \in U : \text{dis}(x, \mathcal{M}) < a\}$  is included in  $\mathcal{V}$ . Let  $\alpha \in A$  and  $x(t)$  be the solution of (6) such that  $x(0) = \alpha$ . Let  $\alpha_0$  be standard in  $\mathcal{V}$  such that  $\alpha \simeq \alpha_0$ . By attractivity of  $\mathcal{M}$ , the solution  $x_0(t)$  of (6) starting by  $\alpha_0$  is defined for all  $t \geq 0$  and satisfies  $\text{dis}(x_0(t), \mathcal{M}) \simeq 0$  for all  $t \simeq +\infty$ . By the Short Shadow Lemma,  $x(t) \simeq x_0(t)$  for all limited  $t > 0$ . By Robinson's Lemma (see for example [17, 13] for this important lemma), there exists  $v \simeq +\infty$  such that  $x(t) \simeq x_0(t)$  for all  $t$  in  $[0, v]$ . Thus,  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all unlimited  $t \leq v$ . By stability of  $\mathcal{M}$ , this approximation still holds for all  $t > v$ . Hence,  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all  $t \geq 0$ . Conversely, if the subset  $\mathcal{M}$  is assumed to satisfy the property in the lemma, then by definition the standard set  $A$  is contained in the basin of attraction of  $\mathcal{M}$ . Hence,  $\mathcal{M}$  is attractive. Let  $x(t)$  be a solution of (6) such that  $x(0) = \alpha$ , where  $\alpha$  is infinitely close to a standard  $\alpha_0 \in \mathcal{M}$ . Since  $\alpha \in A$ , by hypothesis,  $x(t)$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all  $t \simeq +\infty$ . On the other hand, if  $x_0(t)$  is the maximal solution of (6) such that  $x_0(0) = \alpha_0$ , its positive trajectory is in the positively invariant set  $\mathcal{M}$ . Hence, by the Short Shadow Lemma,  $x(t) \simeq x_0(t) \in \mathcal{M}$  for all limited  $t \geq 0$ . Hence,  $\text{dis}(x(t), \mathcal{M}) \simeq 0$  for all  $t \geq 0$  and  $\mathcal{M}$  is stable. ■

**REMARK.** — uniform Let us reconsider the equation (7) depending on the parameter  $y \in Y$ . Suppose that (7) has the property of uniqueness and that  $f$ ,  $U$  and  $Y$  are standard. Let  $\mathcal{M}_y$  be a standard compact subset of  $U$  positively invariant for (7) for all  $y$  in  $Y$ . In the light of the last Lemma, we can also establish that  $\mathcal{M}_y$  is asymptotically stable for the system (7) if and only if there exists a standard  $a > 0$  such that for all standard  $y \in Y$ , any solution  $x(t)$  of (6) for which  $\text{dis}(x(0), \mathcal{M}_y) < a$  can be continued for all  $t \geq 0$  and satisfies  $\text{dis}(x(t), \mathcal{M}_y) \simeq 0$  for all  $t \simeq +\infty$ .

When  $f_0$  and  $g_0$  are standard, we have the following characterization of the SGPAS notion when  $\varepsilon \rightarrow 0$ .

**Lemma 4.4** *Suppose that  $f_0$  and  $g_0$  are standard. A standard bounded subset  $\mathcal{A}$  of  $\Omega_0$  is SGPAS for the system (18) when  $\varepsilon \rightarrow 0$  if and only if for all  $(\alpha, \beta)$  nearstandard in  $\Omega_0$ , all  $\varepsilon > 0$  infinitesimal and all unlimited  $t > 0$ , any solution  $(x(t), y(t))$  of (18) with initial condition  $(\alpha, \beta)$  satisfies  $\text{dis}((x(t), y(t)), \mathcal{A}) \simeq 0$ .*

**Proof.** To say in the Lemma “ $(\alpha, \beta)$  is nearstandard in  $\Omega_0$ ” is the same as to say “ $\exists^{st} \mathcal{K}$ , compact subset of  $\Omega_0$ , such that  $(\alpha, \beta) \in \mathcal{K}$ ”. To say “ $\varepsilon > 0$  infinitesimal” is like to say “ $\forall^{st} \varepsilon_0 > 0, \varepsilon \leq \varepsilon_0$ ”. To say “unlimited  $t > 0$ ” is the same as to say “ $\forall^{st} T t \geq T$ ”. The result  $\text{dis}((x(t), y(t)), \mathcal{A}) \simeq 0$  is equivalent to “ $\forall^{st} \mathcal{O}$  neighborhood of  $\mathcal{A}, (x(t), y(t)) \in \mathcal{O}$ ”. Hence, the characterization of the semiglobal practical asymptotic stability is given in the Lemma by

$$\forall(\alpha, \beta) \forall \varepsilon \forall t (\exists^{st} \mathcal{K} \forall^{st} \varepsilon_0 \forall^{st} T A \Rightarrow \forall^{st} \mathcal{O} (x(t), y(t)) \in \mathcal{O}),$$

where  $A$  is the internal formula  $(\alpha, \beta) \in \mathcal{K}$ ,  $\varepsilon \leq \varepsilon_0$  and  $t \geq T$ . Here,  $f_0, g_0$  and  $\Omega_0$  are standard parameters,  $\varepsilon_0$  and  $T$  positive real numbers and  $\mathcal{K}$  and  $\mathcal{O}$  are subsets. According to the second reduction formula of (19), this is equivalent to

$$\forall \mathcal{K} \forall \mathcal{O} \exists^{fin} \varepsilon'_0 \exists^{fin} T' \\ \forall (\alpha, \beta) \forall \varepsilon \forall t (\forall \varepsilon_0 \in \varepsilon'_0 \forall T \in T' A \Rightarrow (x(t), y(t)) \in \mathcal{O}).$$

The sets  $\varepsilon'_0$  and  $T'$  being finite, there exist  $\varepsilon_0$  and  $T$  such that  $\varepsilon_0 = \min \varepsilon'_0$  and  $T = \max T'$ . The last formula becomes

$$\forall \mathcal{K} \forall \mathcal{O} \exists \varepsilon_0 \exists T \forall (\alpha, \beta) \forall \varepsilon \forall t (A \Rightarrow (x(t), y(t)) \in \mathcal{O}).$$

This is exactly the definition of the semiglobal practical asymptotic stability of  $\mathcal{A}$  for the system (18) when  $\varepsilon \rightarrow 0$ . ■

We can also establish the following characterization which, after all, will be the one we shall use.

**Lemma 4.5** *Suppose that  $f_0$  and  $g_0$  are standard. A standard bounded subset  $\mathcal{A}$  of  $\Omega_0$  is PAS for the system (18) when  $\varepsilon \rightarrow 0$  if and only if there exist a standard compact neighborhood  $\mathcal{K} \subset \Omega_0$  of  $\mathcal{A}$  such that for all  $(\alpha, \beta) \in \mathcal{K}$ , all  $\varepsilon > 0$  infinitesimal and all unlimited  $t > 0$ , any solution  $(x(t), y(t))$  of (18) with initial condition  $(\alpha, \beta)$  satisfies  $dis((x(t), y(t)), \mathcal{A}) \simeq 0$ .*

### 4.3. External notion of perturbation

While the notion of perturbation is classically described via deformations or neighborhoods, a perturbation of a standard object in NSA is just another object, which is nonstandard and infinitely close to it. Its properties are investigated directly without using extra-properties with respect to the parameters of deformation. This way of thinking fits better with the intuitive meaning of the term ‘‘perturbation’’.

**Definition 4.6** *An element  $(\Omega, f, g, \alpha, \beta)$  of  $\mathcal{T}$  is said to be a perturbation of the standard element  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  of  $\mathcal{T}$  if  $\Omega$  contains all the nearstandard elements in  $\Omega_0$ ,  $f(x, y) \simeq f_0(x, y)$  and  $g(x, y) \simeq g_0(x, y)$  for all  $(x, y)$  nearstandard in  $\Omega_0$  and  $\alpha \simeq \alpha_0$ ,  $\beta \simeq \beta_0$ .*

The following lemma makes the link between the above definition of perturbation and the topology of uniform convergence on compacta defined in Section 2 :

**Lemma 4.7** *The element  $(\Omega, f, g, \alpha, \beta)$  of  $\mathcal{T}$  is a perturbation of the standard element  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  of  $\mathcal{T}$  if and only if  $(\Omega, f, g, \alpha, \beta)$  is infinitely close to  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  for the topology of uniform convergence on compacta, that is,  $(\Omega, f, g, \alpha, \beta)$  is in any standard neighborhood of  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ .*

**Proof.** See [13], Lemma 2, page 11. ■

#### 4.4. S-Stability

We need through this subsection to introduce a notion of stability which will lead to results generalizing those of Theorems 3.2 and 3.3. Indeed, they will be expressed, in an external way, in terms of neighborhoods of systems as it was done for approximation results in Section 2. Let the system

$$\dot{x} = f(x), \quad (22)$$

be such that the vector field  $f$  is continuous, not necessarily standard. Suppose for instance that  $f$  is defined on an open subset  $U$  of  $\mathbb{R}^n$ .

**Definition 4.8** A standard bounded subset  $\mathcal{M}$  of  $U$  is said to be *s-globally asymptotically stable (s-GAS)* for the system (22) if for all  $x_0$  nearstandard in  $U$  and all unlimited  $t > 0$ , any solution  $x(t)$  with initial condition  $x_0$  satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$ .

The notions s-GAS and GAS are equivalent if  $f$  is standard<sup>4</sup>(see [11], Proposition 4.1, page 10). For instance, the origin of the system  $\dot{x} = x(\varepsilon x - 1)$ , where  $\varepsilon \simeq 0$ , is s-GAS but not GAS. In the same way, here is the notion of s-(local) asymptotic stability.

**Definition 4.9** A standard bounded subset  $\mathcal{M}$  of  $U$  is said to be *s-asymptotically stable (s-AS)* for the system (22) if there exists a standard compact neighborhood  $\mathcal{K} \subset U$  of  $\mathcal{M}$  such that for all  $x_0 \in \mathcal{K}$  and all unlimited  $t > 0$ , any solution  $x(t)$  with initial condition  $x_0$  satisfies  $\text{dis}(x(t), \mathcal{M}) \simeq 0$ .

REMARK. — s-stab Lemmas 4.4 and 4.5 finally say that if  $f_0$  and  $g_0$  are standard, the subset  $\mathcal{A}$  is SGPAS (respectively PAS) when  $\varepsilon \rightarrow 0$  for the system (18) if and only it is s-GAS (respectively s-AS) for all  $\varepsilon$  infinitesimal.

#### 4.5. External statements

Theorems 4.10 and 4.11 below are the external statements of Theorems 2.3 and 2.4 when  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  is standard. We show for example how Theorem 4.10 reduces by Nelson's algorithm to the internal result of Theorem 2.3.

**Theorem 4.10** Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0) \in \mathcal{T}$  be standard and  $\xi : Y \rightarrow \mathbb{R}^n$  be a standard continuous function. Let hypotheses T1 to T5 be satisfied. Let  $\mathcal{M}$  be a standard closed subset in  $\dot{Y}$  which is positively invariant for the slow equation (9). Suppose that  $\mathcal{M}$  is asymptotically stable for (9) with  $\beta_0$  in its basin of attraction. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (3). Let  $\varepsilon > 0$  be infinitesimal and  $(\Omega, f, g, \alpha, \beta) \in \mathcal{T}$  be a perturbation of  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ . Then any solution  $(x(t), y(t))$  of (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that :

$$\begin{aligned} \varepsilon\omega &\simeq 0, \quad \omega' \simeq +\infty, \\ x(\varepsilon\tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ x(t) &\simeq \xi(\bar{y}(t)) \text{ for } t \in [\varepsilon\omega, \omega'] \\ y(t) &\simeq \bar{y}(t) \text{ for } t \in [0, \omega'], \\ \text{dis}(y(t), \mathcal{M}) &\simeq 0 \text{ for } t \geq \omega', \\ \text{dis}(x(t), \xi(\mathcal{M})) &\simeq 0 \text{ for } t \geq \omega'. \end{aligned} \quad (23)$$

4. In nonstandard analysis, the prefix s (s-properties) permits to distinguish between a standard notion and the corresponding nonstandard one. See for instance [3].

**Proof of Theorem 2.3** Let  $F$  be the formula : “any solution  $(x(t), y(t))$  of (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that (10)” and respectively by  $u_0$  and  $u$  the variables  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$  and  $(\Omega, f, g, \alpha, \beta)$  of  $\mathcal{T}$ . We also design by  $F'$  the formula “any solution  $(x(t), y(t))$  of (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that (23)”. On the other hand, to say that “ $\varepsilon$  is infinitesimal” is the same as to say that “ $\forall^{\text{st}} \varepsilon^*, \varepsilon < \varepsilon^*$ ”, to say that “ $u$  is a perturbation of  $u_0$ ” is the same as to say that “ $u$  is in any standard neighborhood  $\mathcal{V}$  of  $u_0$ ”. Finally, the formula  $F'$  is equivalent to the formula  $\forall^{\text{st}} \eta F$ . Theorem 4.10 is formalized by

$$\forall \varepsilon \forall u (\forall^{\text{st}} \varepsilon^* \forall^{\text{st}} \mathcal{V} K \Rightarrow \forall^{\text{st}} \eta F), \tag{24}$$

where  $K$  designates the formula  $\varepsilon < \varepsilon^* \ \& \ u \in \mathcal{V}$ . Here,  $u_0$  is a standard parameter,  $u$  ranges over  $\mathcal{T}$ , while  $\varepsilon$  and  $\varepsilon^*$  range over the positive real numbers and  $\mathcal{V}$  ranges over the neighborhoods of  $u_0$ . Using the first reduction formula of (19), (24) is equivalent to

$$\forall \eta \exists^{\text{fin}} \varepsilon^{*'} \exists^{\text{fin}} \mathcal{V}' \forall \varepsilon \forall u (\forall \varepsilon^* \in \varepsilon^{*''} \forall \mathcal{V} \in \mathcal{V}' K \Rightarrow F).$$

The sets  $\varepsilon^{*'}$  and  $\mathcal{V}'$  being finite, there exist  $\varepsilon^*$  and  $\mathcal{V}$  such that  $\varepsilon^* = \min \varepsilon^{*'}$  and  $\mathcal{V} = \bigcap_{\mathcal{V}' \in \mathcal{V}'} \mathcal{V}'$ . The last formula becomes equivalent to

$$\forall \eta \exists \varepsilon^* \exists \mathcal{V} \forall \varepsilon \forall u (K \Rightarrow F).$$

Hence, the statement of Theorem 2.3 holds for any standard  $u_0$ . By transfer, it holds for any  $u_0$  in  $\mathcal{T}$ .

**Theorem 4.11** Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0) \in \mathcal{T}$  be standard. Let hypotheses P1 to P5 be satisfied. Let  $\mathcal{M}$  be a standard closed subset in  $\hat{G}$  which is positively invariant for the slow equation (11). Suppose that  $\mathcal{M}$  is asymptotically stable for (11) with  $\beta_0$  in its basin of attraction. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (5). Let  $\varepsilon > 0$  be infinitesimal and  $(\Omega, f, g, \alpha, \beta) \in \mathcal{T}$  be a perturbation of  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ . Then any solution  $(x(t), y(t))$  of (1) is defined for all  $t \geq 0$  and there exist  $\omega > 0$  and  $\omega' > 0$  such that :

$$\begin{aligned} \varepsilon \omega &\simeq 0, \ \omega' \simeq +\infty, \\ x(\varepsilon \tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ y(t) &\simeq \bar{y}(t) \text{ for } t \in [0, \omega'], \\ \text{dis}(x(t), \Gamma_{\bar{y}(t)}) &\simeq 0 \text{ for } t \in [\varepsilon \omega, \omega'], \\ \text{dis}(y(t), \mathcal{M}) &\simeq 0 \text{ for } t \geq \omega', \\ \text{dis}(x(t), \Gamma_{y(t)}) &\simeq 0 \text{ for } t \geq \omega'. \end{aligned} \tag{25}$$

According to of section 4.4, the external statements of Theorems 3.2 and 3.3 are stated as follows :

**Theorem 4.12** Let  $f_0 : \Omega_0 \rightarrow \mathbb{R}^n$  and  $g_0 : \Omega_0 \rightarrow \mathbb{R}^m$  be standard continuous functions defined on an open subset  $\Omega_0$  of  $\mathbb{R}^{n+m}$  and satisfying hypotheses T1 to T5. Let  $\mathcal{M}$  be a standard closed subset in  $\hat{Y}$  which is positively invariant for the slow equation (9). Suppose that  $\mathcal{M}$  is asymptotically stable for (9). Then the subset  $\xi(\mathcal{M}) \times \mathcal{M}$  of the slow manifold is s-AS for the system (18) for all  $\varepsilon > 0$  infinitesimal.

**Theorem 4.13** Let  $f_0 : \Omega_0 \rightarrow \mathbb{R}^n$  and  $g_0 : \Omega_0 \rightarrow \mathbb{R}^m$  be standard continuous functions defined on an open subset  $\Omega_0$  of  $\mathbb{R}^{n+m}$  and satisfying hypotheses P1 to P5. Let  $\mathcal{M}$  be a standard closed subset in  $\mathring{G}$  which is positively invariant for the slow equation (11). Suppose that  $\mathcal{M}$  is asymptotically stable for (11). Then the subset  $\bigcup_{y \in \mathcal{M}} (\Gamma_y \times \{y\})$  of  $\Omega_0$  is s-AS for the system (18) for all  $\varepsilon > 0$  infinitesimal.

To prove the internal theorems is by this way equivalent than to prove their external formulations. However, we rather prove two results that contain those of the two last theorems. Let us introduce the set

$$\mathcal{U} = \{(\Omega, f, g) : \Omega \text{ open subset of } \mathbb{R}^{n+m}, \\ f : \Omega \rightarrow \mathbb{R}^n, g : \Omega \rightarrow \mathbb{R}^m \text{ continuous}\}$$

for which we define the notion of perturbation of a standard element  $(\Omega_0, f_0, g_0)$  analogously as in Definition 4.6. Consider the slow and fast system

$$\begin{aligned} \varepsilon \dot{x} &= f(x, y), \\ \dot{y} &= g(x, y), \end{aligned} \quad (26)$$

corresponding to an element  $(\Omega, f, g)$  of  $\mathcal{U}$ .

**Theorem 4.14** Let  $(\Omega_0, f_0, g_0)$  be a standard element of  $\mathcal{U}$  satisfying hypotheses T1 to T5. Suppose that the slow equation (9) has in  $\mathring{Y}$  a closed standard positively invariant subset  $\mathcal{M}$  which is asymptotically stable. Then, the subset  $\xi(\mathcal{M}) \times \mathcal{M}$  of the slow manifold is s-AS for the system (26) for all perturbation  $(\Omega, f, g) \in \mathcal{U}$  of  $(\Omega_0, f_0, g_0)$  and for all  $\varepsilon > 0$  infinitesimal.

**Theorem 4.15** Let  $(\Omega_0, f_0, g_0)$  be a standard element of  $\mathcal{U}$  satisfying hypotheses P1 to P5. Suppose that the slow equation (11) has in  $\mathring{G}$  a closed standard positively invariant subset  $\mathcal{M}$  which is asymptotically stable. Then, the subset  $\bigcup_{y \in \mathcal{M}} (\Gamma_y \times \{y\})$  of  $\Omega_0$  is s-AS for the system (26) for all perturbation  $(\Omega, f, g) \in \mathcal{U}$  of  $(\Omega_0, f_0, g_0)$  and for all  $\varepsilon > 0$  infinitesimal.

## 4.6. Some required theorems and lemmas

Lobry, Sari and Touhami [13] extended Tykhonov's Theorem on compact time interval to systems belonging to a small neighborhood of the unperturbed problem as follows :

**Theorem 4.16** [13] Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0) \in \mathcal{T}$  be standard and  $\xi : Y \rightarrow \mathbb{R}^n$  a continuous standard function. Let hypotheses T1 to T5 be satisfied. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (3). Let  $T$  be a standard real number in the positive interval of definition of  $\bar{y}(t)$ . Let  $\varepsilon > 0$  be infinitesimal and  $(\Omega, f, g, \alpha, \beta) \in \mathcal{T}$  be a perturbation of  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ . Then, any solution  $(x(t), y(t))$  of the problem (1) is defined at least on  $[0, T]$  and there exists a real number  $\omega > 0$  (independent of  $T$ ) such that :

$$\begin{aligned} \varepsilon \omega &\simeq 0, \\ x(\varepsilon \tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ x(t) &\simeq \xi(\bar{y}(t)) \text{ for } \varepsilon \omega \leq t \leq T, \\ y(t) &\simeq \bar{y}(t) \text{ for } 0 \leq t \leq T. \end{aligned}$$

In [20], Sari and Yadi gave the following topological formulation of Pontryagin’s Theorem on finite interval of time :

**Theorem 4.17** *Let  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0) \in \mathcal{T}$  be standard. Let hypotheses P1 to P4 be satisfied. Let  $\tilde{x}(\tau)$  and  $\bar{y}(t)$  be the respective solutions of the boundary layer equation (4) and of the reduced problem (5). Let  $L$  be a standard real number in the positive interval of definition of  $\bar{y}(t)$ . Let  $\varepsilon > 0$  be infinitesimal and  $(\Omega, f, g, \alpha, \beta) \in \mathcal{T}$  be a perturbation of  $(\Omega_0, f_0, g_0, \alpha_0, \beta_0)$ . Then, any solution  $(x(t), y(t))$  of the problem (1) is defined at least on  $[0, L]$  and there exists a real number  $\omega > 0$  (independent of  $L$ ) such that :*

$$\begin{aligned} \varepsilon\omega &\simeq 0, \\ x(\varepsilon\tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ y(t) &\simeq \bar{y}(t) \text{ for } 0 \leq t \leq L, \\ x(t) &\simeq \Gamma_{\bar{y}(t)} \text{ for } \varepsilon\omega \leq t \leq L. \end{aligned}$$

The proof of Theorem 4.10 needs two facts established in [13]. The first is a lemma which states that a trajectory of the original problem (1) which comes infinitely close to the slow manifold remains infinitely close to it as long as the  $y$ -component is not close to the boundary of the compact subset  $Y$ . The second lemma says that the  $y$ -component of a solution of (1) that is infinitely close to the slow manifold is infinitely close to the solution  $\bar{y}(t)$  of the slow equation (9).

**Lemma 4.18 ([13] Lemma 9, page 17)** *Let hypotheses T1, T2 and T3 be satisfied. Let  $(x(t), y(t))$  be a solution of (1) such that  $y(t)$  is nearstandard in  $\mathring{Y}$  for all  $t$  in  $[t_0, t_1]$  and  $x(t_0) \simeq \xi(y(t_0))$ , then  $x(t) \simeq \xi(y(t))$  for all  $t$  in  $[t_0, t_1]$ .*

**Lemma 4.19 ([13] Lemma 10, page 18)** *Let hypothesis T4 be satisfied. Let  $y_0$  be standard in  $\mathring{Y}$ . Let  $(x(t), y(t))$  be a solution of (1) such that  $x(t) \simeq \xi(y(t))$  for all  $t$  in  $[t_0, t_1]$  and  $y(t_0) \simeq y_0$ . Let  $\bar{y}(t)$  be the solution of the slow equation (9) with initial condition  $y_0$  which is assumed to be defined on the standard interval  $[0, T]$ . Then  $y(t_0 + s) \simeq \bar{y}(s)$  for all  $s \leq T$  such that  $t_0 + s \leq t_1$ .*

The proof of Theorem 4.11 needs two facts established in [20]. The first is a lemma which states that a trajectory of the original problem (1) which comes infinitely close to the manifold generated by the cycles of the fast equation remains infinitely close to it as long as the  $y$ -component is not close to the boundary of the compact subset  $G$ . The second lemma says that the  $y$ -component of a solution of (1) that is infinitely close to this manifold is infinitely close to the solution  $\bar{y}(t)$  of the averaged system (11).

**Lemma 4.20 ([20] Lemma 4.4, page 9)** *Let hypotheses P1 and P2 be satisfied. Let  $(x(t), y(t))$  be a solution of (1) such that  $y(t)$  is nearstandard in  $\mathring{G}$  for  $t \in [t_0, t_1]$  and  $x(t_0) \simeq \Gamma_{y(t_0)}$ . Then  $x(t) \simeq \Gamma_{y(t)}$  for all  $t$  in  $[t_0, t_1]$ .*

**Lemma 4.21 ([20] Lemma 4.5, page 10)** *Let hypotheses P1 to P5 be satisfied. Let  $(x(t), y(t))$  be a solution of (1) such that  $y(t_0)$  is nearstandard in  $\mathring{G}$ . Let  $y_0$  be standard in  $\mathring{G}$  such that  $y(t_0) \simeq y_0$  and  $\bar{y}(t)$  the solution of (11) with initial condition  $y_0$  and defined on the standard interval  $[0, L]$ . Let  $t_1 \geq t_0$  such that  $t_1 \leq t_0 + L$  and  $x(t) \simeq \Gamma_{y(t)}$  for  $t \in [t_0, t_1]$ . Then  $y(t_0 + s) \simeq \bar{y}(s)$  for all  $0 \leq s \leq L$  such that  $t_0 + s \leq t_1$ .*

## 5. Proof of the main results

### 5.1. Proof of Theorem 4.10

Let  $(x(t), y(t))$  be a solution of the problem (1). By assumptions  $T1$  to  $T5$  Theorem 4.16 asserts that there exists a real number  $\omega > 0$  such that :

$$\begin{aligned} \varepsilon\omega &\simeq 0, \\ x(\varepsilon\tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ x(t) &\simeq \xi(\bar{y}(t)) \text{ for } \varepsilon\omega \leq t \leq T, \\ y(t) &\simeq \bar{y}(t) \text{ for } 0 \leq t \leq T, \end{aligned} \quad (27)$$

where  $T$  is a standard real number in the positive interval of definition of the unique solution  $\bar{y}(t)$  of the reduced problem (3). The real number  $\beta_0$  being in the basin of attraction of the subset  $\mathcal{M}$ , the solution  $\bar{y}(t)$  is defined for  $t \geq 0$  and satisfies  $\text{dis}(\bar{y}(t), \mathcal{M}) \simeq 0$  for  $t \simeq +\infty$ . Hence, the approximations (27) hold for any limited real number  $T > 0$ . By Robinson's Lemma, the two last approximations of (27) remain true until a positive unlimited real number  $\omega'$ . This proves the four first approximations of (23). One has  $y(\omega') \simeq \bar{y}(\omega') \simeq \mathcal{M}$  and  $x(\omega') \simeq \xi(\bar{y}(\omega')) \simeq \xi(\mathcal{M})$  by continuity of the standard function  $\xi$ . Now, consider the solution starting at the point  $(x(\omega'), y(\omega'))$ . Let  $y_0(\omega')$  be the standard part of  $y(\omega')$ . It is obvious that  $y_0(\omega') \in \mathcal{M}$ . Since  $\mathcal{M}$  is positively invariant, the positive trajectory of the maximal solution of the slow equation with initial condition  $y_0(\omega')$  lies in  $\mathcal{M}$ . We can apply again Theorem 4.16 to obtain

$$\begin{aligned} y(\omega' + k) &\simeq \mathcal{M}, \\ x(\omega' + k) &\simeq \xi(\mathcal{M}) \text{ for all limited } k \geq 0. \end{aligned} \quad (28)$$

Suppose that there exists  $s \geq \omega'$  such that  $y(s) \not\simeq \mathcal{M}$ , that is there exists a standard positive real number  $\mu$  such that  $\text{dis}(y(s), \mathcal{M}) = \mu$ . One can chose  $s$  in manner to have the set  $\mathcal{A} = \{y \in \mathbb{R}^m : \text{dis}(y, \mathcal{M}) \leq \mu\}$  included in the basin of attraction of  $\mathcal{M}$ . Let  $m$  be the smallest value of  $s$  which exists by compactness of the boundary of  $\mathcal{A}$ . Thus,  $\text{dis}(y(m), \mathcal{M}) = \mu$ . The component  $y(t)$  is then nearstandard in  $\dot{Y}$  for all  $t$  in  $[\omega', m]$  and  $x(\omega') \simeq \xi(y(\omega'))$ . By Lemma 4.18,  $x(t) \simeq \xi(y(t))$  for all  $t$  in  $[\omega', m]$ . On the other hand, the time  $k_0 := m - \omega'$  needed to go from  $(x(\omega'), y(\omega'))$  to  $(x(m), y(m))$  is positive unlimited. Indeed, if  $k_0$  was limited, by (28) one should have  $y(\omega' + k_0) \simeq \mathcal{M}$ , that is  $y(m) \simeq \mathcal{M}$  which contradicts the fact that  $\text{dis}(y(m), \mathcal{M}) = \mu$ . Now, the solution starting from  $(x(m), y(m))$  satisfies  $y(m + k) \in \mathcal{A}$  for all  $k$  in  $[-k_0, 0]$ . Let  $\bar{y}(k)$  the solution of the slow equation (9) with initial condition  $\bar{y}(0) = y_0(m)$ , where  $y_0(m)$  is the standard part of  $y(m)$ . Lemma 4.19 ensures that  $y(m + k) \simeq \bar{y}(k)$  for all limited  $k \leq 0$ . According to Robinson's lemma, there exists  $k_1 < 0$  unlimited such that  $y(m + k_1) \simeq \bar{y}(k_1)$ . One can chose  $k_1$  such that  $-k_0 \leq k_1$ . Hence,  $\bar{y}(k_1)$  is in  $\mathcal{A}$ , thus in the basin of attraction of  $\mathcal{M}$ , and, according to Lemma 4.3, the asymptotic stability of  $\mathcal{M}$  gives  $\bar{y}(k_1 + k) \simeq \mathcal{M}$  for all  $k > 0$  unlimited. In particular, for  $k = -k_1$ , one has  $\bar{y}(0) \simeq \mathcal{M}$ . But  $\bar{y}(0) = y_0(m)$  and  $y_0(m) \simeq y(m)$ , thus  $y(m) \simeq \mathcal{M}$ . This is a contradiction with the definition of  $m$ . Finally, (28) is satisfied for all  $k \geq 0$  which proves the two last approximations of (23).

### 5.2. Proof of Theorem 4.11

Let  $(x(t), y(t))$  be a solution of the problem (1). By assumptions  $P1$  to  $P5$ , Theorem 4.17 asserts that there exists a real number  $\omega > 0$  such that :

$$\begin{aligned} \varepsilon\omega &\simeq 0, \\ x(\varepsilon\tau) &\simeq \tilde{x}(\tau) \text{ for } 0 \leq \tau \leq \omega, \\ y(t) &\simeq \bar{y}(t) \text{ for } 0 \leq t \leq L, \\ x(t) &\simeq \Gamma_{\bar{y}(t)} \text{ for } \varepsilon\omega \leq t \leq L, \end{aligned} \tag{29}$$

for all limited  $L > 0$ . The number  $\beta_0$  being in the basin of attraction of the subset  $\mathcal{M}$ , the solution  $\bar{y}(t)$  is defined for  $t \geq 0$  and satisfies  $\text{dis}(\bar{y}(t), \mathcal{M}) \simeq 0$  for  $t \simeq +\infty$ . Hence, the approximations (29) hold for any limited real number  $L > 0$ . By Robinson's Lemma, the two last approximations of (29) remain true until a positive unlimited real number  $\omega'$ . This proves the four first approximations of (25). One has  $y(\omega') \simeq \bar{y}(\omega') \simeq \mathcal{M}$  and  $x(\omega') \simeq \Gamma_{\bar{y}(\omega')} \simeq \Gamma_{y(\omega')}$  by the continuous dependence of the orbits  $\Gamma_y$  with respect to  $y$ . As in the preceding proof, the fact that  $\mathcal{M} \subset \overset{\circ}{G}$  is positively invariant and the application of Theorem 4.17 to the solution starting at the point  $(x(\omega'), y(\omega'))$  lead to the approximations

$$\begin{aligned} y(\omega' + k) &\simeq \mathcal{M}, \\ x(\omega' + k) &\simeq \Gamma_{y(\omega'+k)} \text{ for all limited } k \geq 0. \end{aligned} \tag{30}$$

Suppose that there exists  $s \geq \omega'$  such that  $y(s) \not\simeq \mathcal{M}$ , that is there exists a standard positive real number  $\mu$  such that  $\text{dis}(y(s), \mathcal{M}) = \mu$ . One can chose  $s$  in manner to have the set  $\mathcal{A} = \{y \in \mathbb{R}^m : \text{dis}(y, \mathcal{M}) \leq \mu\}$  included in the basin of attraction of  $\mathcal{M}$ . Let  $m$  be the smallest value of  $s$  which exists by compactness of the boundary of  $\mathcal{A}$ . Thus,  $\text{dis}(y(m), \mathcal{M}) = \mu$ . The component  $y(t)$  is then nearstandard in  $\overset{\circ}{G}$  for all  $t$  in  $[\omega', m]$  and  $x(\omega') \simeq \Gamma_{y(\omega')}$ . By Lemma 4.20,  $x(t) \simeq \Gamma_{y(t)}$  for all  $t$  in  $[\omega', m]$ . Here again, the time  $k_0 := m - \omega'$  needed to go from  $(x(\omega'), y(\omega'))$  to  $(x(m), y(m))$  is positive unlimited. Now, the solution starting from  $(x(m), y(m))$  satisfies  $y(m+k) \in \mathcal{A}$  for all  $k$  in  $[-k_0, 0]$ . Let  $\bar{y}(k)$  the solution of the slow equation (11) with initial condition  $\bar{y}(0) = y_0(m)$ , where  $y_0(m)$  is the standard part of  $y(m)$ . Lemma 4.21 ensures that  $y(m+k) \simeq \bar{y}(k)$  for all limited  $k \leq 0$ . According to Robinson's lemma, there exists  $k_1 < 0$  unlimited such that  $y(m+k_1) \simeq \bar{y}(k_1)$ . One can chose  $k_1$  such that  $-k_0 \leq k_1$ . Hence,  $\bar{y}(k_1)$  is in  $\mathcal{A}$ , thus in the basin of attraction of  $\mathcal{M}$ , and, according to Lemma 4.3, the asymptotic stability of  $\mathcal{M}$  gives  $\bar{y}(k_1+k) \simeq \mathcal{M}$  for all  $k > 0$  unlimited. In particular, for  $k = -k_1$ , one has  $\bar{y}(0) \simeq \mathcal{M}$ . But  $\bar{y}(0) = y_0(m)$  and  $y_0(m) \simeq y(m)$ , thus  $y(m) \simeq \mathcal{M}$ . This is a contradiction with the definition of  $m$ . Finally, (30) is satisfied for all  $k \geq 0$  which proves the two last approximations of (25).

### 5.3. Proofs of Theorems 4.14 and 4.15

Let  $a > 0$  be the uniform bound deduced from Remark ???. Let  $b > 0$  be a standard real number such that the compact neighborhood  $\{y \in \mathbb{R}^m : \text{dis}(y, \mathcal{M}) \leq b\}$  of  $\mathcal{M}$  is in the basin of attraction of  $\mathcal{M}$ . Then consider the compact neighborhood of  $\xi(\mathcal{M}) \times \mathcal{M}$  on the slow manifold defined by

$$\mathcal{K} = \{(x, y) \in \Omega_0 : \|x - \xi(y)\| \leq a/2, \text{dis}(y, \mathcal{M}) \leq b\}.$$

Let  $(\Omega, f, g)$  be a perturbation of  $(\Omega_0, f_0, g_0)$  and  $\varepsilon > 0$  infinitesimal. Since  $\Omega$  contains all the nearstandard points in  $\Omega_0$ , we have  $\mathcal{K} \subset \Omega$ . Let  $(\alpha, \beta)$  be an arbitrarily point of  $\mathcal{K}$ . We want to show that any solution  $(x(t), y(t))$  of (26) is defined for all  $t \geq 0$  and satisfies  $(x(t), y(t)) \simeq \xi(\mathcal{M}) \times \mathcal{M}$  for all  $t \simeq +\infty$ . Let  $(\alpha_0, \beta_0)$  be standard such that  $(\alpha, \beta) \simeq (\alpha_0, \beta_0)$ . By compactness, we have  $(\alpha_0, \beta_0) \in \mathcal{K}$ . Theorem 4.10 and the construction of  $\mathcal{K}$  imply that for all  $\varepsilon \simeq 0$ , any solution  $(x(t), y(t))$  of (26) is defined for all  $t \geq 0$  and there exists  $\omega' \simeq +\infty$  such that

$$\begin{aligned} y(t) &\simeq \bar{y}(t) \text{ for } 0 \leq t \leq \omega', \\ x(t) &\simeq \xi(\bar{y}(t)) \text{ for } 0 < t \leq \omega', \\ \text{dis}(y(t), \mathcal{M}) &\simeq 0 \text{ for } t \geq \omega', \\ \text{dis}(x(t), \xi(\mathcal{M})) &\simeq 0 \text{ for } t \geq \omega', \end{aligned} \tag{31}$$

where  $\bar{y}(t)$  is the solution of the reduced problem (5). Hence, for  $t \geq \omega'$ , we already have  $(x(t), y(t)) \simeq \xi(\mathcal{M}) \times \mathcal{M}$ . On the other hand, since  $\text{dis}(\beta_0, \mathcal{M}) < b$ , the asymptotic stability of  $\mathcal{M}$  implies that  $\text{dis}(\bar{y}(t), \mathcal{M}) \simeq 0$  for all  $t \simeq +\infty$ . In particular, this is true for all unlimited  $0 < t \leq \omega'$ . According to the first and second approximations of (31),  $\text{dis}(y(t), \mathcal{M}) \simeq \text{dis}(\bar{y}(t), \mathcal{M}) \simeq 0$  and  $\text{dis}(x(t), \xi(\mathcal{M})) \simeq 0$  for all unlimited  $0 < t \leq \omega'$ . Theorem 4.14 is proved, knowing that  $(\Omega, f, g)$  was an arbitrary perturbation of  $(\Omega_0, f_0, g_0)$ .

The proof of Theorem 4.15 can be done in the same way as above. It suffices to consider the compact neighborhood

$$\mathcal{K} = \{(x, y) \in \Omega_0 : \text{dis}(x, \Gamma_y) \leq a, \text{dis}(y, \mathcal{M}) \leq b\}$$

of the set  $\bigcup_{y \in \mathcal{M}} (\Gamma_y \times \{y\})$  where  $a$  is given by Remark ?? and  $b$  is a standard real number such that the set  $\{y \in \mathbb{R}^m : \text{dis}(y, \mathcal{M}) \leq b\}$  is in the basin of  $\mathcal{M}$ . For any  $(\alpha, \beta) \in \mathcal{K}$ , Theorem 4.11 leads to the result.

#### 5.4. Proofs of Theorems 4.12 and 4.13

These Theorems are obviously particular cases of Theorems 4.14 and 4.15 where  $(\Omega_0, f_0, g_0)$  is not perturbed. Note that the proof of Theorems 4.14 and 4.15 show how stability results are deduced from the same assumptions as those of Theorems 4.10 and 4.11.

**Acknowledgement** This work was partially done during my numerous stays in Mulhouse University (France) with the help of my advisor Pr Tewfik SARI. I would like to thank him as well as the referee for valuable comments and suggestions.

---

## 6. References

- [1] D.V. ANOSOV, «On limit cycles in systems of differential equations with a small parameter in the highest derivatives», *AMS Translations*, Ser. 2 33 (1963) 233-275.
- [2] J.L. CALLOT, T. SARI, «Stroboscopie et moyennisation dans les systèmes d'équations différentielles à solutions rapidement oscillantes», *Mathematical Tools and Models for Control, Systems Analysis and Signal Processing*, 3, CNRS Paris (1983) 345-353.

- [3] F. DIENER, M. DIENER (Eds), «*Nonstandard Analysis in Practice*», Universitext Springer-Verlag, 1995.
- [4] M. FARKAS, «*Periodic Motions*», Springer-Verlag, Applied Mathematical Sciences 104, 1994.
- [5] N. FENICHEL, «Geometric singular perturbation theory for ordinary differential equations», *J. Diff. Eq.*, 31 (1979) 53-98.
- [6] L. FLATTO, N. LEVINSON, «Periodic solutions of singularly perturbed systems», *J. Rational Mech.*, 4 (1955) 943-950.
- [7] K. O. FRIEDRICHS, W. WASOW, «Singular perturbations of nonlinear oscillations», *Duke Math. J.* 13 (1946) 361-381.
- [8] W. HAHN, «*Stability of Motion*», Springer-Verlag, Heidelberg, 1967.
- [9] H. K. KHALIL, «*Nonlinear Systems*», Prentice Hall, 1996.
- [10] M. LAKRIB, «Time averaging for functional differential equations», *J. Appl. Math.* 2003, 1 (2003) 1-16.
- [11] C. LOBRY, T. SARI, «The Peaking Phenomenon and Singular Perturbations : An Extension of Tikhonov's Theorem», Research Report n° 4051 of Institut National de Recherche en Informatique et Automatique, 2000.
- [12] C. LOBRY, T. SARI, «Singular perturbations methods in control theory», in *Contrôle Non Linéaire et Applications, Cours du CIMPA, Collection Travaux en Cours*, Hermann éditeur, (2005) 151-177.
- [13] C. LOBRY, T. SARI, S. TOUHAMI, «On Tykhonov's theorem for convergence of solutions of slow and fast systems», *Electron. J. Diff. Eqns*, 19 (1998) 1-22.
- [14] R. LUTZ, «L'intrusion de l'analyse non standard dans l'étude des perturbations singulières», *Astérisque* 109-110, (1983) 101-140.
- [15] E. NELSON, «Internal Set Theory : a new approach to nonstandard analysis», *Bull. Amer. Math. Soc.*, 83 (1977) 1165-1198.
- [16] L. S. PONTRYAGIN, L. V. RODYGIN, «Approximate solution of a system of ordinary differential equations involving a small parameter in the derivatives», *Soviet. Math. Dokl.*, 1 (1960) 237-240.
- [17] A. ROBINSON, «*Nonstandard Analysis*», American Elsevier, New York, 1974.
- [18] T. SARI, «Petite histoire de la stroboscopie», In *Colloque Trajectorien à la mémoire de Georges Reeb et Jean-louis Callot*, ed. by A. Fruchard and A. Troesh (IRMA Publication, 1995) 5-15.
- [19] T. SARI, «Stroboscopy and averaging», In *Colloque Trajectorien à la mémoire de Georges Reeb et Jean-louis Callot*, ed. by A. Fruchard and A. Troesh (IRMA Publication, 1995) 95-124.
- [20] T. SARI, K. YADI, «On Pontryagin–Rodygin's theorem for convergence of solutions of slow and fast systems», *Electron. J. Diff. Eqns*, 19 (2004), 1-17.
- [21] A. R. TEEL, L. PRALY, «Tools for semiglobal stabilization by partial state and output feedback», *SIAM J. Contr. Optim.* 33 (1995) 1443-1485.
- [22] A. N. TYKHONOV, «Systems of differential equations containing small parameters multiplying the derivatives», *Mat. Sborn.*, 31 (1952) 575-586.
- [23] F. VERHULST, «Periodic solutions and slow manifolds», *Int. J. Bifurc. Chaos Appl. Sci. Engrg* 17 (2007) 2533-2540.
- [24] F. VERHULST, T. BAKRI, «The dynamics of slow manifold», *J. Indones. Math. Soc.* **13** (2007) n°1, 73-90.
- [25] A. K. ZVONKIN, M. A. SHUBIN, «Nonstandard analysis and singular perturbations of ordinary differential equations», *Uspehi Mat. Nauk.* 39 (1984) 77-127.